

# Local Tait colourings of cubic graphs

Martin Škoviera

**Comenius University, Bratislava**

Bordeaux Graph Workshop, 18-20 November, 2010

**In honour of André Raspaud**

# Contents

- **Part I:** Local Tait colourings
  - generalisation of 3-edge-colourings
  - relation to Steiner triple systems and point-line configurations
  - existence of local Tait colourings
- **Part II:** Local Tait colourings of bridgeless cubic graphs
  - Fano colourings
  - abelian colourings
  - integer  $k$ -colourings
  - colourings by symmetric configurations
- **Conclusion**
  - Hierarchy of conjectures related to local Tait colourings


## Credits

D. Archdeacon, H.-L. Fu, M. Grannell, T. Griggs, F. Holroyd, M. Knor, D. Král',  
E. Máčajová, O. Pangrác, A. Pór, A. Raspaud, J.-S. Sereni, & M. S.

# Edge-colourings of cubic graphs

*Edge-colouring* of a graph ... assignment of colours to its edges such that adjacent edges receive distinct colours

- First studied by P. G. Tait (1880)
- J. Petersen (1898) ... four colours suffice to colour every cubic (= trivalent) graph



# Edge-colourings of cubic graphs

*Chromatic index*  $\chi'(G)$  ... minimum # of colours needed to colour  $G$

*Class 1* ... graphs with  $\chi' = 3$  (Tait-colourable graphs)

*Class 2* ... graphs with  $\chi' = 4$

# Edge-colourings of cubic graphs

*Chromatic index*  $\chi'(G)$  ... minimum # of colours needed to colour  $G$

*Class 1* ... graphs with  $\chi' = 3$  (Tait-colourable graphs)

*Class 2* ... graphs with  $\chi' = 4$

Well-known:

- colouring by 4 colours ... *easy*
- deciding whether  $\chi' = 3$  or 4 ... *difficult* ... Holyer (1981)
- almost all cubic graphs are *Class 1* ... Wormald (1992)

# Edge-colourings of cubic graphs

*Chromatic index*  $\chi'(G)$  ... minimum # of colours needed to colour  $G$

*Class 1* ... graphs with  $\chi' = 3$  (Tait-colourable graphs)

*Class 2* ... graphs with  $\chi' = 4$

Well-known:

- colouring by 4 colours ... *easy*
- deciding whether  $\chi' = 3$  or 4 ... *difficult* ... Holyer (1981)
- almost all cubic graphs are *Class 1* ... Wormald (1992)

Non-trivial graphs with  $\chi' = 4$  (called *snarks*) closely related to

- Cycle-Double Cover Conj., Fulkerson's Conj., 5-Flow Conj., etc.

# Generalisation of 3-edge-colouring

## Local Tait colourings

- allow an arbitrary number of colours
- **global** condition on  $\#$  of colours  $\rightarrow$  **local** condition:

(L) *any two colours meeting at a vertex always determine the same the third colour*

# Generalisation of 3-edge-colouring

## Local Tait colourings

- allow an arbitrary number of colours
- **global** condition on  $\#$  of colours  $\rightarrow$  **local** condition:

(L) *any two colours meeting at a vertex always determine the same the third colour*

Condition (L) trivially satisfied by usual 3-edge-colourings


$\Rightarrow$  “**local**” Tait colourings generalise “**global**” ones

# Local Tait colourings: example

Consider a **proper** edge-colouring  $\phi : E(G) \rightarrow \mathbb{Z}_3 \times \mathbb{Z}_3$

- colours ... elements of the group  $\mathbb{Z}_3 \times \mathbb{Z}_3$
- colours meeting at a vertex ... triples that sum to **0**

$\Rightarrow \phi$  is a **local** Tait colouring by elements of  $\mathbb{Z}_3 \times \mathbb{Z}_3$



# Geometric viewpoint

$\phi : E(G) \rightarrow P$  ... local Tait colouring with colour set  $P$

- view the colours as geometric points
- place a line through a pair of points whenever the corresponding colours meet at a vertex

# Geometric viewpoint

$\phi : E(G) \rightarrow P$  ... local Tait colouring with colour set  $P$

- view the colours as geometric **points**
- place a **line** through a pair of points whenever the corresponding colours meet at a vertex

Then:

- By Condition (L), there is **at most one line** through a pair of points
- Each line contains **exactly three points** corresponding to colours

# Geometric viewpoint

$\phi : E(G) \rightarrow P \dots$  local Tait colouring with colour set  $P$

- view the colours as geometric **points**
- place a **line** through a pair of points whenever the corresponding colours meet at a vertex

Then:

- By Condition (L), there is **at most one line** through a pair of points
- Each line contains **exactly three points** corresponding to colours

Every local Tait colouring determines a **configuration of points and lines**.

# Combinatorial viewpoint

$\phi : E(G) \rightarrow P \dots$  local Tait colouring with

- $P \dots$  set of colours
- $B \dots$  triples of colours occurring at vertices

# Combinatorial viewpoint

$\phi : E(G) \rightarrow P \dots$  local Tait colouring with

- $P \dots$  set of colours
- $B \dots$  triples of colours occurring at vertices

By Condition (L):

*any two elements of  $P$  belong to **at most** one triple listed in  $B$*

$\Rightarrow$  the pair  $(P, B)$  forms a **partial Steiner triple system**

# Steiner triple systems & configurations

A *Steiner triple system*  $\mathcal{S} = (P, B)$  of order  $n$  consists of

- set  $P$  of  $n$  points
- collection  $B$  of 3-element subsets of  $P$  (called *blocks*) s.t. any two points belong to **exactly** one block

If any two points belong to **at most one** one block

⇒ *partial Steiner triple system*

# Steiner triple systems & configurations

A *Steiner triple system*  $\mathcal{S} = (P, B)$  of order  $n$  consists of

- set  $P$  of  $n$  points
- collection  $B$  of 3-element subsets of  $P$  (called *blocks*) s.t. any two points belong to **exactly** one block

If any two points belong to **at most one** one block

⇒ *partial Steiner triple system*

Well-known:

- Any partial STS embeds into some full STS ... [Treash \(1976\)](#)  
(may need some extra vertices)
- ⇒ Any partial STS can be viewed as a *configuration* of points and blocks of a full STS

# Colourings by configurations

Summing up:

Any local Tait colouring can be viewed as one where

- colours ... **points** of a (partial) Steiner triple system  $\mathcal{S}$
- colours that meet at a vertex form a **block** of  $\mathcal{S}$

$\Rightarrow$   *$\mathcal{S}$ -colouring*

# Colourings by configurations

Summing up:

Any local Tait colouring can be viewed as one where

- colours ... **points** of a (partial) Steiner triple system  $\mathcal{S}$
- colours that meet at a vertex form a **block** of  $\mathcal{S}$

$\Rightarrow$   $\mathcal{S}$ -colouring

Tait colouring =  $\mathcal{I}$ -colouring,  $\mathcal{I}$  being the trivial STS of order 3

# Projective and affine STS's

The *projective* Steiner triple system  $PG(n, 2)$ ,  $n \geq 2$ , has

- points ...  $P = \mathbb{Z}_2^{n+1} - \{0\}$
- blocks ... triples  $\{x, y, z\}$  with  $x + y + z = 0$

smallest projective STS ...  $PG(2, 2)$  ... **Fano plane** of order **7**

# Projective and affine STS's

The *projective* Steiner triple system  $PG(n, 2)$ ,  $n \geq 2$ , has


- points ...  $P = \mathbb{Z}_2^{n+1} - \{0\}$
- blocks ... triples  $\{x, y, z\}$  with  $x + y + z = 0$

smallest projective STS ...  $PG(2, 2)$  ... **Fano plane** of order **7**

The *affine* Steiner triple system  $AG(n, 3)$ ,  $n \geq 2$ , has

- points ...  $P = \mathbb{Z}_3^n$
- blocks ... triples  $\{x, y, z\}$  with  $x + y + z = 0$

smallest affine STS ...  $AG(2, 3)$  ... **affine plane** of order **9**

Fano plane  $PG(2, 2)$ , the smallest non-trivial STS

# Existence of $\mathcal{S}$ -colourings: bridgeless graphs

**Problem** (Archdeacon, 1986). Is it possible to find an  $\mathcal{S}$ -colouring of a cubic graph for a given Steiner triple system  $\mathcal{S}$  ?

# Existence of $\mathcal{S}$ -colourings: bridgeless graphs

**Problem** (Archdeacon, 1986). Is it possible to find an  $\mathcal{S}$ -colouring of a cubic graph for a given Steiner triple system  $\mathcal{S}$  ?

- (Fu, 2001): Bridgeless cubic graphs of **genus**  $\leq 24$  or of **order**  $\leq 189$  are **Fano**-colourable

## Existence of $\mathcal{S}$ -colourings: bridgeless graphs

**Problem** (Archdeacon, 1986). Is it possible to find an  $\mathcal{S}$ -colouring of a cubic graph for a given Steiner triple system  $\mathcal{S}$  ?


- (Fu, 2001): Bridgeless cubic graphs of **genus**  $\leq 24$  or of **order**  $\leq 189$  are **Fano**-colourable

**Theorem** (Holroyd & S., 2004)

*Every bridgeless cubic graph has an  $\mathcal{S}$ -colouring for every non-trivial Steiner triple system  $\mathcal{S}$ .*


# Existence of $\mathcal{S}$ -colourings: graphs with bridges

There exist graphs with **no** local Tait colouring!




Existence of  $\mathcal{S}$ -colourings: graphs with bridges

There exist graphs with **no** local Tait colouring!




Existence of  $\mathcal{S}$ -colourings: graphs with bridges

There exist graphs with **no** local Tait colouring!





Existence of  $\mathcal{S}$ -colourings: graphs with bridges

There exist graphs with **no** local Tait colouring!



Existence of  $\mathcal{S}$ -colourings: graphs with bridges

There exist graphs with **no** local Tait colouring!



*series-parallel* = no subdivision of  $K_4$

# Existence of $\mathcal{S}$ -colourings: graphs with bridges

- Every cubic graph with **no series-parallel end** admits a local **Tait** colouring.

# Existence of $\mathcal{S}$ -colourings: graphs with bridges

- Every cubic graph with **no series-parallel end** admits a local **Tait** colouring.

Proof. Take  $(P, B)$  with

$P = E(G)$  and  $B =$  triples of pairwise adjacent edges.

# Existence of $\mathcal{S}$ -colourings: graphs with bridges

- Every cubic graph with **no series-parallel end** admits a local **Tait** colouring.

Proof. Take  $(P, B)$  with

$P = E(G)$  and  $B =$  triples of pairwise adjacent edges.

- Question: Does there exist a *universal* STS?

(STS is *universal*  $\Leftrightarrow$  it colours all **simple** cubic graphs)

# Existence of $\mathcal{S}$ -colourings: graphs with bridges

- Every cubic graph with **no series-parallel end** admits a local **Tait** colouring.

Proof. Take  $(P, B)$  with

$P = E(G)$  and  $B =$  triples of pairwise adjacent edges.

- Question: Does there exist a *universal* STS?

(STS is *universal*  $\Leftrightarrow$  it colours all **simple** cubic graphs)

- Projective and affine systems are **not** universal!

Projective systems do not colour **graphs with bridges**

Affine systems do not colour graphs with **bipartite ends**

# Universal Steiner triple systems

- Universal STS of order 381 (Grannell, Griggs, Knor, S., 2004)

# Universal Steiner triple systems

- Universal STS of order 381 (Grannell, Griggs, Knor, S., 2004)
- Universal point-transitive STS of order 21 minimal under inclusion, (a subsystem of the 381-system) (Pál & S., 2007)

# Universal Steiner triple systems

- Universal STS of order 381 (Grannell, Griggs, Knor, S., 2004)
- Universal point-transitive STS of order 21 minimal under inclusion, (a subsystem of the 381-system) (Pál & S., 2007)
- Every non-projective and non-affine point-transitive STS is universal.  
⇒ The smallest order of a universal STS is 13.  
(Kráľ', Máčajová, Pór, Sereni, 2010)

# Universal Steiner triple systems

- Universal STS of order 381 (Grannell, Griggs, Knor, S., 2004)
- Universal point-transitive STS of order 21 minimal under inclusion, (a subsystem of the 381-system) (Pál & S., 2007)
- Every non-projective and non-affine point-transitive STS is universal.  
⇒ The smallest order of a universal STS is 13.  
(Kráľ', Máčajová, Pór, Sereni, 2010)

Question: Point-intransitive systems?

# Universal Steiner triple systems

Universality of **point-intransitive** STS's  
(Grannell, Griggs, Máčajová, S., 2010+):

- Infinitely many point-intransitive inclusion-minimal **universal** systems  
(**Wilson-Schreiber systems** = abelian group + two extra points)
- Infinitely many point-intransitive systems that are **not universal**  
(projective systems  $PG(2, n)$ ,  $n \geq 3$ , modified by one **Pasch switch**)

## PART II

# Local Tait colourings of bridgeless cubic graphs

# Colourings by configurations in STS's

## Theorem (Holroyd & S., 2004)

*Every bridgeless cubic graph has an  $\mathcal{S}$ -colouring for every non-trivial Steiner triple system  $\mathcal{S}$ .*


# Colourings by configurations in STS's

## Theorem (Holroyd & S., 2004)

*Every bridgeless cubic graph has an  $\mathcal{S}$ -colouring for every non-trivial Steiner triple system  $\mathcal{S}$ .*

Question: What happens if a Steiner triple system  $\mathcal{S}$  is replaced by a suitable configuration?

# 1. Fano colourings



**Fano colouring** – proper edge-colouring of a cubic graph

- **colours** – points of the Fano plane
- the colours around each vertex **form a line**

# 1. Fano colourings

How many **points/lines** of the **Fano plane** are needed to colour a given cubic graph  $G$ ?

# 1. Fano colourings

How many **points/lines** of the **Fano plane** are needed to colour a given cubic graph  $G$ ?

- If  $G$  is 3-edge-colourable
  - any **single line** is sufficient.

# 1. Fano colourings

How many **points/lines** of the **Fano plane** are needed to colour a given cubic graph  $G$ ?


- If  $G$  is 3-edge-colourable
  - any **single line** is sufficient.
- If  $G$  is not 3-edge-colourable
  - all **seven points**
  - at least **four lines** are needed.

# 1. Fano colourings

How many **points/lines** of the **Fano plane** are needed to colour a given cubic graph  $G$ ?

- If  $G$  is 3-edge-colourable
  - any **single line** is sufficient.
- If  $G$  is not 3-edge-colourable
  - all **seven points**
  - at least **four lines** are needed.

**$F_i$ -colouring** – colouring using at most  $i$  lines of the Fano plane

$F_5$ -colouring of the Petersen graph

# $F_7$ -colourings

All seven lines are never needed:

# $F_i$ -colourings

All seven lines are never needed:

Theorem (Máčajová & S., 2005)

$F_6$ T

*Every bridgeless cubic graph admits an  $F_6$ -colouring.*

$F_i$ -colourings

All seven lines are never needed:

Theorem (Máčajová & S., 2005)

 $F_6T$ 

*Every bridgeless cubic graph admits an  $F_6$ -colouring.*

We believe that four lines are always enough:

4-Line Conjecture (Máčajová & S., 2005)

 $F_4C$ 

*Every bridgeless cubic graph admits an  $F_4$ -colouring.*

# Conjecture of Fan and Raspaud

## Theorem (Máčajová & S., 2005)

The *4-Line Conjecture* is equivalent to the following conjecture of Fan and Raspaud:

## Conjecture (Fan & Raspaud, 1994)

F&amp;RC

Every bridgeless cubic graphs contains **three** perfect matchings with **no edge in common**.

# Conjecture of Fan and Raspaud – background

Theorem (Petersen, 1891)

*Every bridgeless cubic graphs contains a perfect matching.*

# Conjecture of Fan and Raspaud – background

## Theorem (Petersen, 1891)

*Every bridgeless cubic graphs contains a perfect matching.*

- A cubic graphs has **two disjoint** perfect matchings  
 $\Leftrightarrow$  **3**-edge-colourable.

# Conjecture of Fan and Raspaud – background

## Theorem (Petersen, 1891)

*Every bridgeless cubic graphs contains a perfect matching.*

- A cubic graphs has **two disjoint** perfect matchings  
 $\Leftrightarrow$  **3**-edge-colourable.
- Every **two** perfect matchings in a non-**3**-edge-colourable graph have an **edge in common**.

# Conjecture of Fan and Raspaud – background

Conjecture (Fan & Raspaud, 1994)

F&RC

Every bridgeless cubic graph contains **three** perfect matchings with **no edge in common**.

# Conjecture of Fan and Raspaud – background

## Conjecture (Fan & Raspaud, 1994)

F&amp;RC

Every bridgeless cubic graphs contains **three** perfect matchings with **no edge in common**.

## Conjecture (Berge, Fulkerson, 1971)


FC

Every bridgeless cubic graphs contains a family of **six** perfect matchings which together cover each edge exactly twice.

# Conjecture of Fan and Raspaud – background

**Definition.** A configuration  $\mathcal{C}$  is *Class 1* if  $\mathcal{C} \rightarrow \mathcal{I}$ , otherwise it is *Class 2*.


**Example:** The smallest *Class 2* configuration is  $C_{15} \cong F_4$ , the *sail*.



# Conjecture of Fan and Raspaud – background

**Definition.** A configuration  $\mathcal{C}$  is *Class 1* if  $\mathcal{C} \rightarrow \mathcal{I}$ , otherwise it is *Class 2*.

**Example:** The smallest *Class 2* configuration is  $C_{15} \cong F_4$ , the *sail*.



## 4-Line-Conjecture (rephrased)

The smallest *Class 2* configuration colours every bridgeless cubic graph.

# Conjecture of Fan and Raspaud

$$\mathbf{FC} \Rightarrow \mathbf{F\&RC} \Leftrightarrow F_4C \Rightarrow F_5C \Rightarrow F_6T \equiv \mathbf{TRUE}$$

# Conjecture of Fan and Raspaud

$$\mathbf{FC} \Rightarrow \mathbf{F\&RC} \Leftrightarrow F_4C \Rightarrow F_5C \Rightarrow F_6T \equiv \mathbf{TRUE}$$

Theorem (Kaiser & Raspaud, 2007, 2010)

*Every bridgeless cubic graph of oddness  $\leq 2$  admits an  $F_5$ -colouring.*

oddness – minimum number of odd circuits in a 2-factor  
 oddness 0  $\Leftrightarrow$  Class 1

# Conjecture of Fan and Raspaud

$$\mathbf{FC} \Rightarrow \mathbf{F\&RC} \Leftrightarrow F_4C \Rightarrow F_5C \Rightarrow F_6T \equiv \mathbf{TRUE}$$

Theorem (Kaiser & Raspaud, 2007, 2010)

*Every bridgeless cubic graph of oddness  $\leq 2$  admits an  $F_5$ -colouring.*

oddness – minimum number of odd circuits in a 2-factor  
 oddness 0  $\Leftrightarrow$  Class 1

Theorem (Máčajová & S., 2009+)

*Every bridgeless cubic graph of oddness  $\leq 2$  admits an  $F_4$ -colouring.*

Equivalently:

*Every bridgeless cubic graph of oddness  $\leq 2$  has **three** perfect matchings with **no edge in common**.*

## 2. Abelian colourings

- Given an abelian group  $A$ , an  $A$ -colouring of a cubic graph is a proper colouring by elements of  $A - 0$  such that around each vertex the colours sum to zero.

## 2. Abelian colourings

- Given an abelian group  $A$ , an  $A$ -colouring of a cubic graph is a proper colouring by elements of  $A - 0$  such that around each vertex the colours sum to zero.

Abelian colourings are local Tait colourings:

- Let  $A$  be an abelian group. Define  $\mathcal{C}(A)$  to be the partial STS where
  - points ...  $A - 0$
  - blocks ... triples  $\{x, y, z\}$  with  $x + y + z = 0$

Clearly,  $A$ -colouring =  $\mathcal{C}(A)$ -colouring

## 2. Abelian colourings

- Given an abelian group  $A$ , an  $A$ -colouring of a cubic graph is a proper colouring by elements of  $A - 0$  such that around each vertex the colours sum to zero.

Abelian colourings are local Tait colourings:

- Let  $A$  be an abelian group. Define  $\mathcal{C}(A)$  to be the partial STS where
  - points ...  $A - 0$
  - blocks ... triples  $\{x, y, z\}$  with  $x + y + z = 0$

Clearly,  $A$ -colouring =  $\mathcal{C}(A)$ -colouring

### Remark.

$A$ -colouring  $\leftrightarrow$  n.-z.  $A$ -flow on a cubic graph with *antibalanced bidirection*

## 2. Abelian colourings

- If  $A$  is one of  $\mathbb{Z}_2$ ,  $\mathbb{Z}_3$ ,  $\mathbb{Z}_4$ , and  $\mathbb{Z}_5$ , then  $\mathcal{C}(A) = \emptyset \Rightarrow$  no  $A$ -colouring
- If  $A$  is one of  $\mathbb{Z}_2 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_6$ ,  $\mathbb{Z}_7$ ,  $\mathbb{Z}_8$ , and  $\mathbb{Z}_9$ , then  
 $A$ -colouring  $\Leftrightarrow$  3-edge-colouring

## 2. Abelian colourings

- If  $A$  is one of  $\mathbb{Z}_2$ ,  $\mathbb{Z}_3$ ,  $\mathbb{Z}_4$ , and  $\mathbb{Z}_5$ , then  $\mathcal{C}(A) = \emptyset \Rightarrow$  no  $A$ -colouring
- If  $A$  is one of  $\mathbb{Z}_2 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_6$ ,  $\mathbb{Z}_7$ ,  $\mathbb{Z}_8$ , and  $\mathbb{Z}_9$ , then  
 $A$ -colouring  $\Leftrightarrow$  3-edge-colouring

Theorem (Máčajová, Raspaud & S., 2005)

If  $A = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$  or  $|A| \geq 12 \Rightarrow$  every bridgeless cubic graph is  $A$ -colourable.

## 2. Abelian colourings


- If  $A$  is one of  $\mathbb{Z}_2$ ,  $\mathbb{Z}_3$ ,  $\mathbb{Z}_4$ , and  $\mathbb{Z}_5$ , then  $\mathcal{C}(A) = \emptyset \Rightarrow$  no  $A$ -colouring
- If  $A$  is one of  $\mathbb{Z}_2 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_6$ ,  $\mathbb{Z}_7$ ,  $\mathbb{Z}_8$ , and  $\mathbb{Z}_9$ , then  
 $A$ -colouring  $\Leftrightarrow$  3-edge-colouring

Theorem (Máčajová, Raspaud & S., 2005)


If  $A = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$  or  $|A| \geq 12 \Rightarrow$  every bridgeless cubic graph is  $A$ -colourable.

Proof.


Each of  $F_7$ ,  $D_9$ , or  $D_8$  colours all bridgeless cubic graphs (Holroyd & S.):



$F_7$



$D_9$



$D_8$

# The four remaining groups

## Remaining groups:

$\mathbb{Z}_4 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_3 \times \mathbb{Z}_3$ ,  $\mathbb{Z}_{10}$ , and  $\mathbb{Z}_{11} \Rightarrow ???$

# The four remaining groups

## Remaining groups:

$\mathbb{Z}_4 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_3 \times \mathbb{Z}_3$ ,  $\mathbb{Z}_{10}$ , and  $\mathbb{Z}_{11} \Rightarrow ???$

- For each  $A \in \{\mathbb{Z}_4 \times \mathbb{Z}_2, \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_{10}, \mathbb{Z}_{11}\}$  we have  $F_4 \subseteq \mathcal{C}(A)$   
 $\Rightarrow$  4-Line-Conjecture implies  $A$ -colouring

# The four remaining groups

## Remaining groups:

$\mathbb{Z}_4 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_3 \times \mathbb{Z}_3$ ,  $\mathbb{Z}_{10}$ , and  $\mathbb{Z}_{11} \Rightarrow ???$

- For each  $A \in \{\mathbb{Z}_4 \times \mathbb{Z}_2, \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_{10}, \mathbb{Z}_{11}\}$  we have  $F_4 \subseteq \mathcal{C}(A)$   
 $\Rightarrow$  4-Line-Conjecture implies  $A$ -colouring
- $\mathcal{C}(\mathbb{Z}_4 \times \mathbb{Z}_2) = F_5$

# The four remaining groups

## Remaining groups:

$\mathbb{Z}_4 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_3 \times \mathbb{Z}_3$ ,  $\mathbb{Z}_{10}$ , and  $\mathbb{Z}_{11} \Rightarrow ???$

- For each  $A \in \{\mathbb{Z}_4 \times \mathbb{Z}_2, \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_{10}, \mathbb{Z}_{11}\}$  we have  $F_4 \subseteq \mathcal{C}(A)$   
 $\Rightarrow$  4-Line-Conjecture implies  $A$ -colouring
- $\mathcal{C}(\mathbb{Z}_4 \times \mathbb{Z}_2) = F_5$
- $F_5 \not\subseteq \mathcal{C}(A)$  for the other three groups

# The four remaining groups

## Remaining groups:

$\mathbb{Z}_4 \times \mathbb{Z}_2$ ,  $\mathbb{Z}_3 \times \mathbb{Z}_3$ ,  $\mathbb{Z}_{10}$ , and  $\mathbb{Z}_{11} \Rightarrow ???$

- For each  $A \in \{\mathbb{Z}_4 \times \mathbb{Z}_2, \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_{10}, \mathbb{Z}_{11}\}$  we have  $F_4 \subseteq \mathcal{C}(A)$   
 $\Rightarrow$  4-Line-Conjecture implies  $A$ -colouring
- $\mathcal{C}(\mathbb{Z}_4 \times \mathbb{Z}_2) = F_5$
- $F_5 \not\subseteq \mathcal{C}(A)$  for the other three groups







Theorem (Kráľ', Máčajová, Pangrác, Raspaud, Sereni & S., 2008)

*The 5-Line Conjecture implies the existence of an  $A$ -colouring of every bridgeless cubic graph for each  $A \in \{\mathbb{Z}_4 \times \mathbb{Z}_2, \mathbb{Z}_3 \times \mathbb{Z}_3, \mathbb{Z}_{10}, \mathbb{Z}_{11}\}$*

## Equivalence of colourings by Fano plane and by affine plane

Theorem (Kráľ, Máčajová, Pangrác, Raspaud, Sereni & S., 2008)

There exist  $i$ -line configurations  $A_i \subseteq AG(2, 3)$  and  $F_i \subseteq PG(2, 2)$ ,  $4 \leq i \leq 6$ , s. t. a cubic graph has an  $F_i$ -colouring  $\Leftrightarrow$  has an  $A_i$ -colouring.



### 3. Integer colourings

- An *integer  $k$ -colouring* of a cubic graph is a proper  $\mathbb{Z}$ -colouring  $\sigma$  satisfying the condition  $0 < |\sigma(e)| < k$  for each edge  $e$ .

### 3. Integer colourings

- An *integer  $k$ -colouring* of a cubic graph is a proper  $\mathbb{Z}$ -colouring  $\sigma$  satisfying the condition  $0 < |\sigma(e)| < k$  for each edge  $e$ .

Integer  $k$ -colourings are **local Tait colourings**:

- Define  $\mathcal{I}_k$  to be the partial STS where
  - points ... all integers  $n$  with  $|n| < k$
  - blocks ... triples  $\{x, y, z\}$  with  $x + y + z = 0$

Clearly, integer  $k$ -colouring =  $\mathcal{I}_k$ -colouring

### 3. Integer colourings

Theorem (Kráľ', Máčajová, Pangrác, Raspaud, Sereni & S., 2008)

Let  $G$  be a bridgeless cubic graph.

(1) If  $G$  admits an  $F_i$ -colouring,  $i \in \{4, 5, 6\}$ , then it also admits an integer  $(i + 2)$ -colouring.

### 3. Integer colourings

Theorem (Kráľ', Máčajová, Pangrác, Raspaud, Sereni & S., 2008)

Let  $G$  be a bridgeless cubic graph.

- (1) If  $G$  admits an  $F_i$ -colouring,  $i \in \{4, 5, 6\}$ , then it also admits an integer  $(i + 2)$ -colouring.
- (2) If  $G$  admits an integer 6-colouring, then it admits both a  $\mathbb{Z}_{10}$ -colouring and a  $\mathbb{Z}_{11}$ -colouring.

### 3. Integer colourings

Theorem (Kráľ, Máčajová, Pangrác, Raspaud, Sereni & S., 2008)

Let  $G$  be a bridgeless cubic graph.

- (1) If  $G$  admits an  $F_i$ -colouring,  $i \in \{4, 5, 6\}$ , then it also admits an integer  $(i + 2)$ -colouring.
- (2) If  $G$  admits an integer 6-colouring, then it admits both a  $\mathbb{Z}_{10}$ -colouring and a  $\mathbb{Z}_{11}$ -colouring.

#### Corollary

Every bridgeless cubic graph has an integer 8-colouring.

### 3. Integer colourings

Theorem (Kráľ, Máčajová, Pangrác, Raspaud, Sereni & S., 2008)

Let  $G$  be a bridgeless cubic graph.

- (1) If  $G$  admits an  $F_i$ -colouring,  $i \in \{4, 5, 6\}$ , then it also admits an integer  $(i + 2)$ -colouring.
- (2) If  $G$  admits an integer 6-colouring, then it admits both a  $\mathbb{Z}_{10}$ -colouring and a  $\mathbb{Z}_{11}$ -colouring.

#### Corollary

Every bridgeless cubic graph has an integer 8-colouring.

#### Conjecture

Every bridgeless cubic graph admits an integer 6-colouring.

## 4. Colourings by symmetric configurations

**Definition.** *Symmetric configuration*  $n_3$  ... partial STS which has

- $n$  points and  $n$  blocks
- each point contained in exactly three blocks

## 4. Colourings by symmetric configurations

**Definition.** *Symmetric configuration*  $n_3$  ... partial STS which has

- $n$  points and  $n$  blocks
- each point contained in exactly three blocks

**Examples:**


$7_3$  = Fano plane (Class 2)

$8_3$  = Moebius-Kantor configuration =  $AG(2, 3)$ -pt (Class 2)

$9_3$  = Pappus configuration (Class 1)

$10_3$  = Desargues configuration (Class 2)

$15_3$  = Cremona-Richmond configuration (Class 2)


Desargues configuration  $10_3$ 

# Desargues colouring and 5-CDC

**5-Cycle-Double-Cover Conjecture.** *Every bridgeless graph contains a collection of  $\leq 5$  cycles (even subgraphs) s.t. each edge belongs to exactly two of them.*

## Theorem

*A cubic graph has a Desargues colouring  $\Leftrightarrow$  it has a 5-CDC.*

Cremona-Richmond configuration  $15_3$ 

# Cremona-Richmond colouring and Fulkerson's conjecture

**Fulkerson's Conjecture.** *Every bridgeless **cubic** graph contains a collection of **six** perfect matchings s.t. each edge is contained in exactly **two** of them.*

## Theorem

*A cubic graph has a **Cremona-Richmond colouring**  $\Leftrightarrow$  it has a **double covering** of its edges by **six** perfect matchings.*

# Cremona-Richmond configuration and Petersen colourings

## Petersen colouring conjecture

The edges of every bridgeless cubic graph  $G$  can be mapped into the edges of the **Petersen graph** in such a way that any three mutually incident edges of  $G$  are mapped to three mutually incident edges of the **Petersen graph**.

**Definition.** Such a mapping is called a *Petersen colouring* of  $G$ .

# Cremona-Richmond configuration and Petersen colourings

## Petersen colouring conjecture

The edges of every bridgeless cubic graph  $G$  can be mapped into the edges of the **Petersen graph** in such a way that any three mutually incident edges of  $G$  are mapped to three mutually incident edges of the **Petersen graph**.

**Definition.** Such a mapping is called a *Petersen colouring* of  $G$ .

- A Petersen colouring is a **local Tait colouring**.

Define  $\mathcal{P}$  to be the partial STS where

points ... edges of the Petersen graph

blocks ... triples  $\{x, y, z\}$  incident to the same vertex

# Cremona-Richmond configuration and Petersen colourings

## Petersen colouring conjecture

The edges of every bridgeless cubic graph  $G$  can be mapped into the edges of the **Petersen graph** in such a way that any three mutually incident edges of  $G$  are mapped to three mutually incident edges of the **Petersen graph**.

**Definition.** Such a mapping is called a *Petersen colouring* of  $G$ .

- A Petersen colouring is a **local Tait colouring**.

Define  $\mathcal{P}$  to be the partial STS where

points ... edges of the Petersen graph

blocks ... triples  $\{x, y, z\}$  incident to the same vertex


$\mathcal{P} =$  *depleted Cremona-Richmond configuration*

$= 15_3 - \{\text{parallel class of blocks}\}.$

# PART III

## Conclusion

## III. Conclusion: hierarchy of conjectures



**THANK YOU!**